

表 3 不同提取条件下清蛋白浓度($\mu\text{g}/\mu\text{L}$)

Table 3 Albumin concentration under different extraction conditions ($\mu\text{g}/\mu\text{L}$)

提取次数 Times	A	B	C	D	E	F	G	H
1	1.24	1.45	2.12	1.47	1.87	2.06	1.55	1.46
2	1.46	0.90	0.62	0.53	0.61	0.67	0.69	0.59
3	0.53	0.46	0.35	0.31	0.38	0.32	0.41	0.36
平均值 AVG	1.08	0.94	1.03	0.77	0.95	1.02	0.88	0.83

注: A,B,C,D: 分别按 30 $\mu\text{L}/\text{mg}$, 40 $\mu\text{L}/\text{mg}$, 50 $\mu\text{L}/\text{mg}$, 70 $\mu\text{L}/\text{mg}$ 加入提取液, 振荡 10 min; E,F,G,H: 分别按 50 $\mu\text{L}/\text{mg}$ 加入提取液, 振荡 5 min, 15 min, 30 min, 45 min

Note: A,B,C,D: Separately according to 30 $\mu\text{L}/\text{mg}$, 40 $\mu\text{L}/\text{mg}$, 50 $\mu\text{L}/\text{mg}$ and 70 $\mu\text{L}/\text{mg}$ join extract, shock 10 min; E,F,G,H: Separately according to 50 $\mu\text{L}/\text{mg}$ join extract, shock 5 min, 15 min, 30 min and 45 min