

表1南宁市北郊农耕区冬季鸟类组成

Table 1 Winter bird constitute in northern part cultivated land of Nanning

| 种类 Species | Pi (%) | | | | | 居留型 Resident |
|-------------------------------|----------------|-----------------------|------------------------|---------------------------|-----------------------------|-----------------|
| | 湿地 Wet land | 干稻田 Dry rice field | 蔬菜地 Vegetable field | 荒草地 Wild grass f field | 稀树灌丛 Less-tree bush wood | |
| 牛背鹭 | 0.077 9 | 0.060 8 | | | | R |
| <i>Bubulcus ibis</i> | | | | | | |
| 池鹭 | 0.129 9 | | | | | R |
| <i>Ardeola bacchus</i> | | | | | | |
| 绿鹭 | 0.013 0 | | | | | R |
| <i>Butorides striatus</i> | | | | | | |
| 黑翅鸢 | | 0.008 0 | | | 0.011 4 | R |
| <i>Elanus caeruleus</i> | | | | | | |
| 红隼 | | 0.009 6 | | | 0.005 7 | R |
| <i>Falco tinnunculus</i> | | | | | | |
| 白胸苦恶鸟 | 0.051 9 | | 0.005 5 | | | R |
| <i>Amaurornis phoenicurus</i> | | | | | | |
| 金眶鸻 | 0.116 9 | | | | | W |
| <i>Charadrius dubius</i> | | | | | | |
| 扇尾沙锥 | 0.129 9 | 0.016 0 | | | | W |
| <i>Gallinago gallinago</i> | | | | | | |
| 白腰草鹬 | 0.026 0 | | | | | W |
| <i>Tringa ochropus</i> | | | | | | |
| 珠颈斑鸠 | | 0.019 2 | 0.054 6 | | 0.011 4 | R |
| <i>Streptopelia chinensis</i> | | | | | | |
| 褐翅鸦鹃 | | | | | 0.008 5 | R |
| <i>Centropus sinensis</i> | | | | | | |
| 普通翠鸟 | 0.064 9 | | | | | R |
| <i>Alcedo atthis</i> | | | | | | |
| 白胸翡翠 | 0.013 0 | | 0.005 5 | | 0.002 8 | R |
| <i>Halcyon smyrnensis</i> | | | | | | |
| 白鹡鸰 | 0.129 9 | 0.060 8 | 0.207 7 | 0.075 2 | | R |
| <i>Motacilla alba</i> | | | | | | |
| 黄鹡鸰 | 0.1558 | 0.0288 | | 0.112 8 | | W |
| <i>Motacilla flava</i> | | | | | | |
| 灰鹡鸰 | 0.039 0 | | | 0.015 0 | | W |
| <i>Motacilla cinerea</i> | | | | | | |
| 田鹨 | 0.013 0 | 0.043 2 | 0.010 9 | 0.052 6 | | W |
| <i>Anthus richardi</i> | | | | | | |
| 树鹨 | 0.026 0 | 0.160 0 | 0.0273 | 0.210 5 | 0.017 0 | W |
| <i>Anthus hodgsoni</i> | | | | | | |
| 红耳鹎 | | 0.009 6 | 0.043 7 | | 0.156 3 | R |
| <i>Pycnonotus jocosus</i> | | | | | | |
| 白头鹎 | | | | | 0.065 3 | R |
| <i>Pycnonotus sinensis</i> | | | | | | |
| 白喉红臀鹎 | | | | | 0.085 2 | R |
| <i>Pycnonotus aurigaster</i> | | | | | | |
| 红尾伯劳 | | 0.001 6 | 0.005 5 | | 0.005 7 | W |
| <i>Lanius cristatus</i> | | | | | | |
| 棕背伯劳 | | 0.032 0 | 0.005 5 | 0.015 0 | 0.019 9 | R |
| <i>Lanius schach</i> | | | | | | |

续表

Continuing table

| 种类 Species | <i>Pi</i> (%) | | | | | 居留型 Resident |
|----------------------------------|----------------|-----------------------|------------------------|-------------------------|-----------------------------|-----------------|
| | 湿地 Wet land | 干稻田 Dry rice field | 蔬菜地 Vegetable field | 荒草地 Wild grass field | 稀树灌丛 Less-tree bush wood | |
| | | | | | | |
| 八哥 | 0.012 8 | 0.043 7 | | 0.011 4 | | R |
| <i>Acridotheres cristatellus</i> | | | | | | |
| 黑领椋鸟 | 0.009 6 | 0.043 7 | | 0.011 4 | | R |
| <i>Sturnus nigricollis</i> | | | | | | |
| 灰椋鸟 | 0.009 6 | | | 0.073 9 | | W |
| <i>Sturnus cineraceus</i> | | | | | | |
| 灰燕鵙 | 0.006 4 | | | | | R |
| <i>Artamus fuscus</i> | | | | | | |
| 蓝喉歌鸲 | 0.013 0 | | | | | W |
| <i>Luscinia svecicus</i> | | | | | | |
| 鹟 | | 0.010 9 | | 0.034 1 | | R |
| <i>Copsychus saularis</i> | | | | | | |
| 黑喉石 | 0.033 6 | 0.098 4 | 0.060 2 | 0.008 5 | | W |
| <i>Saxicola torquata</i> | | | | | | |
| 灰林 | 0.025 6 | 0.054 6 | 0.090 2 | 0.022 7 | | R |
| <i>Saxicola ferrea</i> | | | | | | |
| 乌 | | | | 0.011 4 | | R |
| <i>Turdus merula</i> | | | | | | |
| 棕扇尾 | 0.001 6 | | | | | R |
| <i>Cisticola juncidis</i> | | | | | | |
| 黄腹山 | | | 0.015 0 | 0.017 0 | | R |
| <i>Prinia flaviventris</i> | | | | | | |
| 纯色山 | 0.028 8 | 0.043 7 | 0.180 5 | 0.022 7 | | R |
| <i>Prinia subflava</i> | | | | | | |
| 长尾缝叶 | | | 0.007 5 | 0.062 5 | | R |
| <i>Orthotomus sutorius</i> | | | | | | |
| 褐柳 | | | | 0.011 4 | | W |
| <i>Phylloscopus fuscatus</i> | | | | 0.039 8 | | R |
| 暗绿绣眼 | | | | | | |
| <i>Zosterops japonica</i> | | | | | | |
| 大山雀 | | | | 0.034 1 | | R |
| <i>Parus major</i> | | | | | | |
| 麻雀 | 0.304 0 | 0.229 5 | 0.150 4 | 0.170 5 | | R |
| <i>Passer montanus</i> | | | | | | |
| 白腰文 | 0.016 0 | | | 0.039 8 | | R |
| <i>Lonchura striata</i> | | | | | | |
| 斑文 | 0.048 0 | | | | | R |
| <i>Lonchura punctulata</i> | | | | | | |
| 金翅雀 | 0.012 8 | 0.065 6 | | 0.011 4 | | R |
| <i>Carduelis sinica</i> | | | | | | |
| 凤头 | 0.019 2 | 0.010 9 | | 0.017 0 | | R |
| <i>Carduelis sinica</i> | | | | | | |
| 小 | 0.022 4 | 0.032 8 | 0.015 0 | 0.011 4 | | W |
| <i>Emberiza pusilla</i> | | | | | | |

注: *pi*为在该生境种出现个体数占总数的百分比, 无数据表格表示该生境无统计对象

Note: *pi* represent for the species occur in the percentage of the total number of individuals, empty table means no statistics objective