

表 2 各材料不同处理的成苗率和干物重

Table 2 The seedling rate and dry weight of different varieties in different treatments

处理 Treatment	成苗率(%) Seedling rate (%)				干物重(g) Dry weight (g)			
	<i>fst</i>	D5A	黎榆 B LiyuB	滇杂 37 Dianza37	<i>fst</i>	D5A	黎榆 B LiyuB	滇杂 37 Dianza37
稻桩对照 Rice stubble CK	29.5 bB	31.6 cC	25.0 cC	23.5 dBC				
激素处理 Hormone treatment	59.0 aA	62.5 bB	46.4 bB	56.0 bB	7.94 aA	9.82 abA	11.18 aA	8.36 aA
无激素对照 Non hormone CK	43.5 aAB	47.0 bcBC	40.2 bB	37.8 cBC	5.48 aA	6.76 bA	8.09 aA	6.77 aA
种子苗对照 Seeding CK		96.0 aA	93.5 aA	94.0 aA		10.22 aA	10.80 aA	9.92 aA

注: 竖栏中相同字母表示差异不显著; 不同小写字母表示差异达 0.05 显著水平不同大写字母表示差异达 0.01 显著水平

Note: Within the same column, data followed by common letters are not significantly different at 0.05 or 0.01 levels; Data followed by different lowercase letters are significantly different at 0.05 levels; Data followed by different uppercase letters show dramatically significant differences at 0.01 levels