

表 1 1998 年, 2010 年庐山鸟类种群数量及密度

Table 1 Lushan Mountain birds and the density of population in 1998 and 2010

名 称 Name	1998 年(李海涛, 2000) Year 1998		2010 年 Year 2010		比较 1998 年种群密度变化 Comparison of changes in population density in 1998	居留性质 Residential properties
	数量 Number(N)	种群密度 Population density(C)	数量 Number(N)	种群密度 Population density(C)		
鸢 <i>Milvus korschun lineatus</i> (II)	5	0.11	2	0.04	-0.07	R
苍鹰 <i>Accipiter gentils schvedoei</i> (II)	1	0.02	2	0.04	+0.02	T
雀鹰 <i>A. soloensis</i> (II)	1	0.02	3	0.06	+0.04	W
红隼 <i>Falco tinnunculus</i> (II)	2	0.04	5	0.11	+0.07	R
鸺鹠 <i>Glaucidium cuculoides</i> (II)	0	0.00	2	0.04	+0.04	R
灰林鸮 <i>Strix aluco Nivicola</i> (II)	0	0.00	2	0.04	+0.04	R
鹰鹃 <i>Cuculus sparverioides</i>	1	0.02	2	0.04	+0.02	S
四声杜鹃 <i>C. micropterus cropterus</i>	0	0.00	4	0.09	+0.09	R
黑枕绿啄木鸟 <i>Picus canus</i>	0	0.00	12	0.26	+0.26	R
大斑啄木鸟 <i>Dendrocopos major</i>	0	0.00	9	0.19	+0.19	R
戴胜 <i>Upupa epops saturata</i>	0	0.00	4	0.09	+0.09	R
灰鹊鸂 <i>Motacilla cinerea</i>	3	0.06	5	0.11	+0.05	S
黄鹊鸂 <i>M. otacillaflava</i>	0	0.00	8	0.17	+0.17	R
白鹊鸂 <i>M. alba</i>	10	0.21	13	0.28	+0.07	R
树鹊 <i>Anthus hodgsoni</i>	18	0.38	21	0.45	+0.07	R
田鹊 <i>A. rufulus</i>	0	0.00	12	0.26	+0.26	R
山斑鸠 <i>Streptopelia o. orientalis</i>	0	0.00	16	0.34	+0.34	R
长尾山椒鸟 <i>Pericrocotus ethologus</i>	76	1.62	62	1.32	-0.30	R
绿鹦嘴鹎 <i>Spizixos</i>	4	0.09	3	0.06	-0.03	S

<hr/>						
<i>semitorques</i>						
黄臀鹌	57	1.21	42	0.89	-0.32	S
<i>Pycnonotus</i>						
<i>xanthorrhous</i>						
白喉红臀鹌	0	0.00	10	0.21	+0.21	R
<i>Pycnonotus</i>						
<i>aurigaster</i>						
<i>latouchei</i>						
棕背伯劳	5	0.11	6	0.13	+0.02	R
<i>Lanius schach</i>						
虎纹伯劳	1	0.02	0	0.00	-0.02	R
<i>L. tigrinus</i>						
黑卷尾	10	0.21	6	0.13	-0.08	S
<i>Dicrurus</i>						
<i>macrocerus</i>						
灰卷尾	3	0.06	3	0.06	0.00	S
<i>D. leucophaeus</i>						
红嘴蓝鹊	26	0.55	21	0.45	-0.10	R
<i>Cissa</i>						
<i>erythrorhyncha</i>						
大嘴乌鸦	20	0.43	18	0.38	-0.05	R
<i>Corvus</i>						
<i>macrorhynchus</i>						
喜鹊普通亚种	0	0.00	8	0.17	+0.17	R
<i>Pica p. sericea</i>						
金胸歌鸲	1	0.02	3	0.06	+0.04	S
<i>Luscinia</i>						
<i>pectardens</i>						
鹊鸲	13	0.28	22	0.47	+0.19	R
<i>Copsychus</i>						
<i>sauularis</i>						
<i>prosthopellus</i>						
黑喉石鸫	0	0.00	2	0.04	+0.04	R
<i>Saxicola torquata</i>						
<i>rzewalskii</i>						
白腹鸫	1	0.02	0	0.00	-0.02	R
<i>Turdus pallidus</i>						
<i>obseurus</i>						
锈脸钩嘴鹀	13	0.28	21	0.45	+0.17	R
<i>Pomatorhinus</i>						
<i>erythrogenys</i>						
棕颈钩嘴鹀	2	0.04	3	0.06	+0.02	R
<i>P. ruficollis</i>						
小鹪鹩	0	0.00	4	0.09	+0.09	R
<i>p. pusilla</i>						
矛纹草鹀	0	0.00	6	0.13	+0.13	T
<i>Babax</i>						
<i>lanceolatus</i>						
灰翅噪鹀	4	0.09	4	0.09	0.00	W
<i>Garrulax</i>						
<i>cineraceus</i>						
画鹀	3	0.06	2	0.04	+0.02	R
<i>G. c. canorus</i>						
红嘴相思鸟	5	0.11	5	0.11	0.00	R
<i>Liothrix lutea</i>						
褐胁雀鹀	6	0.13	8	0.17	+0.04	R
<i>Alcippe</i>						
<i>ruficapilla</i>						
<hr/>						

<hr/>						
<i>sordidior</i>						
白眶雀鹛	2	0.04	4	0.09	+0.05	S
<i>A. morrisonia</i>						
白领凤鹛	4	0.09	6	0.13	+0.04	R
<i>Yuhina diademata</i>						
黄眉柳莺	7	0.15	10	0.21	+0.06	T
<i>Phylloscopus s.</i>						
<i>subaffinis</i>						
黄腰柳莺	1	0.02	6	0.13	+0.11	W
<i>P.inornatus</i>						
棕腹柳莺	2	0.04	6	0.13	+0.09	R
<i>P. s. subaffinis</i>						
褐头鹟莺	4	0.09	26	0.55	+0.46	R
<i>Prinia subflava</i>						
褐山鹟莺华中亚种	11	0.23	16	0.34	+0.11	R
<i>P. polychroa</i>						
<i>catheria</i>						
方尾鹟	5	0.11	6	0.13	+0.02	S
<i>Culicicapa</i>						
<i>ceylonensis</i>						
<i>calochrysea</i>						
白鹟鹛	4	0.09	8	0.17	+0.08	S
<i>Ficedula</i>						
<i>zanthopygia</i>						
大山雀	78	1.66	66	1.40	-0.22	R
<i>Parus major</i>						
绿背山雀	3	0.06	6	0.13	+0.07	R
<i>P. monticolus</i>						
黄腹山雀	5	0.11	8	0.17	+0.06	R
<i>P. venustus</i>						
红腹山雀	4	0.09	6	0.13	+0.04	R
<i>P. davidi</i>						
红头长尾山雀	55	1.17	42	0.89	-0.28	R
<i>Aegithalos</i>						
<i>concinus</i>						
暗绿锈眼鸟	64	1.36	76	1.62	+0.26	S
<i>Zosterops</i>						
<i>japonica</i>						
灰腹锈眼鸟	43	0.91	59	1.26	+0.35	S
<i>Z. palpebrosa</i>						
<i>siamensis</i>						
麻雀	80	1.70	122	2.60	+0.90	R
<i>Passer montanus</i>						
斑文鸟	11	0.23	9	0.19	-0.04	R
<i>Lonchura</i>						
<i>punctulata</i>						
<i>yunnanensis</i>						
黑头金翅	5	0.11	3	0.06	-0.05	R
<i>Carduelis s.</i>						
<i>sinica</i>						
白斑翅拟蜡嘴雀	5	0.11	6	0.13	+0.02	W
<i>Mycerobas</i>						
<i>melanozanthos</i>						
三道鹟草鹛	9	0.19	11	0.23	+0.04	R
<i>Emberiza cioides</i>						
合计	693	14.74	883	18.79	+4.05	R
Total						
<hr/>						

注: C=N/S, S=47km², “+”表示增加,“-”表示减少;(II)表示国家保护动物等级;“R”表示留鸟,“T”表示旅鸟,“S”表示夏候鸟,“W”表示冬候鸟

Note: C=N/S, S=47km², “+” that the rise, “-” that lower; (II)that the national animal protection grade; “R” that the resident, “T” that the traveler, “S” that the summer migratory birds, “W” that the winter visitor