

表3 不定芽增殖培养培养

Table 3 Proliferation of adventitious buds

处理 Treatment	NAA (mg/L)	6-BA (mg/L)	接种芽数 No. of the buds	增殖频率(%) Frequency of proliferation (%)	平均芽长(厘米) Mean of adventitious buds (cm)	芽基部是否有愈伤组织 Callus on the base of the adventitious buds (Yes or No)
1	0.1	0.5	50	110	5.1	少量愈伤组织 Yes
2	0.1	1	50	168	4.9	少量愈伤组织 Yes
3	0.1	1.5	50	213	4.7	少量愈伤组织, 褐化 Yes, and browning of callus
4	0.5	1	50	312	5	少量愈伤组织, 褐化 Yes, and browning of callus
5	0.5	1.5	50	298	4.8	少量愈伤组织, 褐化 Yes, and browning of callus
6	0.5	2	50	350	4.5	少量愈伤组织, 褐化 Yes, and browning of callus
7	0.5	0.5	50	556	5.7	少量愈伤组织, 褐化 Yes, and browning of callus
8	1	0.5	50	287	4.8	少量愈伤组织, 褐化 Yes, and browning of callus
9	1.5	0.5	50	125	4.9	较多愈伤组织 Yes, and much more

注: 增殖频率=增殖后芽数/接种芽数×100%

Note: Frequency of proliferation equals to percentage of that numbers of proliferated buds divide by numbers of buds