

表 2 山药-苦瓜醇提取物对 2 型糖尿病大鼠血糖的影响(x±s)

Table 2 The effect of the ethanol extract compound on blood sugar in type 2 diabetic rats (x±s)

| 组别 Group category | 例数 Case numbers | 治疗前(mmol/L) Before treatment (mmol/L) | 治疗后(mmol/L) After treatment (mmol/L) |
|--|--------------------|---|--|
| 正常对照 Normal control group | 8 | 3.95±1.02 | 3.99±1.66 |
| 模型对照组 Model control group | 8 | 19.11±2.12 | 19.00±2.57 |
| 山药-苦瓜醇提取物高剂量治疗组 High dose treating group with the alcohol extract compound | 8 | 18.80±2.15Δ | 15.87±1.60 |
| 山药-苦瓜醇提取物中剂量组治疗 Alcohol extract middle-dose group | 8 | 19.96±2.02* | 18.89±2.05 |
| 山药-苦瓜醇提取物低剂量治疗组 Low dose treating group with the alcohol extract compound | 8 | 19.02±2.34 | 18.76±3.34 |
| 山药-苦瓜水提取物高剂量治疗组 High dose treating group with the water extract compound | 8 | 19.47±3.26* | 18.20±2.54 |

注: 前后自身对照; *p<0.05; Δ p<0.01

Note: Before and after the self-control; *p<0.05; Δ p<0.01