

表 2 13 个不同省份间黄连木居群遗传距离

Table 2 Genetic distance of *P. chinensis* populations among 13 provinces

省份 province	湖南 Hunan	河北 Hebei	河南 Henan	云南 Yunnan	安徽 Anhui	山东 Shandong	湖北 Hubei	山西 Shanxi	四川 Sichuan	江苏 Jiangsu	浙江 Zhejiang	甘肃 Gansu
河北 Hebei	0.402 0											
河南 Henan	0.375 1	0.154 0										
云南 Yunnan	0.636 4	0.484 4	0.303 5									
安徽 Anhui	0.758 1	0.286 1	0.375 4	0.319 1								
山东 Shandong	0.903 1	0.258 2	0.364 4	0.501 5	0.126 3							
湖北 Hubei	0.873 2	0.677 7	0.386 1	0.062 0	0.553 2	0.656 8						
山西 Shanxi	0.788 0	0.549 8	0.267 0	0.513 6	0.728 0	0.702 5	0.499 3					
四川 Sichuan	0.546 9	0.495 2	0.402 1	0.644 5	0.481 8	0.685 1	0.840 4	0.724 8				
江苏 Jiangsu	0.947 7	0.720 4	0.597 7	0.533 5	1.018 7	1.211 4	0.688 2	0.894 4	0.454 2			
浙江 Zhejiang	0.422 9	0.258 3	0.367 2	0.391 0	0.427 4	0.496 1	0.684 5	0.898 2	0.580 7	0.743 3		
甘肃 Gansu	0.915 7	0.715 7	0.570 3	0.509 9	0.832 0	1.058 4	0.611 7	0.529 6	0.480 1	0.536 3	0.623 1	
贵州 Guizhou	0.833 0	0.528 4	0.310 1	0.364 5	0.516 7	0.509 9	0.419 0	0.310 1	0.430 1	0.643 5	0.549 4	0.145 9