

表 2 12 对 SSR 标记在不同类型水稻品种中的 Shannon 多样性指数和多态性信息含量

Table 2 Shannon's Information index and polymorphism information content at 12 SSR loci in different types of rice cultivars

标记 Locus	Shannon 多样性指数(I)			多态性信息含量(PIC)		
	Shannon's information index (I)			Polymorphism information content (PIC)		
	籼稻 Indica	粳稻 Japonica	总数 Total	籼稻 Indica	粳稻 Japonica	总数 Total
RM297	1.499 8	1.330 8	1.502 2	0.715 0	0.685 0	0.716 2
RM71	1.313 5	0.708 3	1.279 5	0.650 6	0.345 0	0.632 3
RM85	0.329 8	0.639 0	0.537 8	0.150 4	0.340 0	0.254 8
RM5414	0.797 5	0.981 9	0.820 6	0.491 2	0.525 0	0.495 3
RM274	0.912 5	0.518 2	0.907 9	0.566 2	0.2650	0.560 8
RM190	1.358 8	1.039 7	1.356 3	0.735 8	0.625 0	0.734 9
RM336	1.352 4	1.262 9	1.373 0	0.687 6	0.665 0	0.696 6
RM72	1.274 2	1.319 5	1.293 3	0.681 6	0.715 0	0.687 1
RM219	1.314 0	0.500 4	1.318 3	0.676 8	0.320 0	0.684 6
RM311	0.631 8	0.693 1	0.640 7	0.439 9	0.500 0	0.448 5
RM209	1.421 4	1.345 2	1.427 3	0.716 2	0.730 0	0.720 0
RM19	1.066 4	1.366 2	1.146 4	0.608 7	0.740 0	0.634 4
平均值	1.106 0	0.975 4	1.133 6	0.593 3	0.537 9	0.605 5
mean						