

表1 大麦品种赤霉病抗性评价

Table 1 Evaluation of the resistance to fusarium head blight on barley varieties

品种名称 Name of cultivar	单花滴注病小穗率(%) Proportion of scabbed spikelet (%)			孢子喷雾 Spore spread		抗性等级 Level of resistance
	2011	2010	平均 Average	侵染小穗数 No. infected spikelet	病穗率(%) Proportion of scabbed spike (%)	
苏花 2 号 Suhua 2	2.78	7.64	5.21	1.85	59.09	抗 R
新宇 03~3 Xinyu03~3	3.05	9.28	6.17	2.30	83.33	抗 R
5E003	5.22	8.89	7.06	2.63	100.00	抗 R
苏 B0608 Su B0608	5.63	8.93	7.28	1.11	37.50	抗 R
扬农啤 5 号 Yangnongpi5	4.64	12.14	8.39	2.64	53.85	抗 R
苏啤 3 号 Supi3	4.38	12.95	8.67	2.59	100.00	抗 R
Morrison	5.00	14.11	9.56	1.10	72.72	抗 R
苏啤 4 号 Supi4	5.74	13.56	9.65	2.54	100.00	抗 R
盐 99175 Yan99175	5.92	13.38	9.65	2.00	93.33	中抗 MR
申河 085 Shenhe085	10.17	12.94	11.56	2.81	95.45	中抗 MR
扬辐 6008 Yangfu6008	12.50	12.30	12.40	2.68	100.00	中抗 MR
如东 5218 Rudong5218	12.81	16.18	14.50	2.00	77.78	中抗 MR
连 9723 Lian9723	12.28	18.89	15.59	1.57	100.00	中抗 MR
Cheveron	7.54	26.96	17.25	2.11	100.00	中抗 MR
Vivar	9.86	26.48	18.17	2.80	100.00	中抗 MR
CIho4196	9.02	28.31	18.67	3.60	100.00	中抗 MR
单二 Daner	13.28	26.76	20.02	1.89	56.25	中感 MS
Frederickson	26.32	36.43	31.38	5.10	100.00	中感 MS
甘啤 2 号 Ganpi2	35.82	45.83	40.83	6.90	100.00	感 S