

表 1 不同处理方式下，环己酰亚胺影响萱草萼片/花瓣鲜重比值

Table 1 Different sepal/petal fresh weight ratio under different CHI treatments

时间点 Time	对照 Control	处理花梗 Pedicel treated	均匀涂抹 Tepals treated
7:00	0.83±0.07 aA	0.92±0.1 039 aA	0.83±0.07 cC
9:00	0.9233±0.0 115 aA	0.9 667±0.1 234 aA	1.12±0.0 173 bAB
11:00	0.86±0.01 aA	0.83±0.1 493 aA	1.3433±0.0 351 aA
13:00	0.8 833±0.0 503 aA	0.8 667±0.0 723 aA	1.1 433±0.1 102 bAB
15:00	0.8 367±0.106 aA	0.9±0.0 458 aA	1.0 667±0.1 002 bBC
17:00	0.8 067±0.0802 aA	0.8 767±0.0 503 aA	1.0 533±0.0 503 bBC
19:00	0.7 433±0.1 002 aA	0.81±0.0 819 aA	1.0 167±0.0 416 bcBC

注: ±号后代表标准偏差; 小写字母代表 5% 水平下差异显著性, 大写字母代表 1% 水平下差异显著性

Note: ± represents the standard deviation; lower case letters represent significance at 5% confidence level; capital letters represent significance at 1% confidence level