

表 11 不同处理下萼片中细胞质蔗糖转化酶酶活的变化

Table 11 Changes of cytoplasm invertase activity of sepals under different treatments

时间 Time	对照的萼片 Control sepals	处理花梗的萼片 Sepals from treated pedicel	均匀涂抹的萼片 Sepals treated by CHI	测定方法 Measurement method
7:00	3.2 402±0.1 698 aA	2.6 810±0.0 815 cC	3.2 402±0.1 698 aA	μg/min/mg prot.
12:00	2.1 560±0.1 249 bB	3.4 997±0.0 744 bB	2.0 294±0.0 939 bB	
20:00	2.9 879±0.0 809 aA	4.2 406±0.0 929 aA	1.5 779±0.0 506 cC	
7:00	130.3 533±1.9 100 aA	125.0 064±2.8 887 bB	130.3 533±1.9 100 aA	μg/min/g 鲜重
12:00	85.7 992±2.6 807 bB	140.2 447±3.7 752 cC	92.3 248±3.9 959 bB	
20:00	135.7 666±2.6 071 aA	175.4 482±3.5 988 aA	82.1 999±0.4 810 cC	

注: ±号后代表标准偏差; 小写字母代表 5%水平下差异显著性, 大写字母代表 1%水平下差异显著性

Note: ± represents the standard deviation; lower case letters represent significance at 5% confidence level; capital letters represent significance at 1% confidence level